Draft version February 2, 2008
Preprint typeset using LaTeX style emulateapj v. 11/12/01

arXiv:astro-ph/0305496v1 26 May 2003

UNSTABLE COLD DARK MATTER AND THE CUSPY HALO PROBLEM IN DWARF GALAXIES

F.J. Sánchez-Salcedo
Instituto de Astronomía, UNAM, Ciudad Universitaria, Aptdo. 70 264, C.P. 04510, Mexico City, Mexico
jsanchez@astroscu.unam.mx

Draft version February 2, 2008

ABSTRACT

We speculate that the dark halos of dwarf galaxies and low surface brightness galaxies soften their central cusps by the decay of a fraction of cold dark matter (CDM) particles to a stable particle with a recoiling velocity of a few tens km s^{-1}, after they have driven the formation of galactic halos. This process, however, does not necessarily produce a significant reduction of the central dark matter density of satellite dwarf spheroidals like Draco or Fornax. It is shown that the recovered distribution of concentration parameters c for the initial (before decay) Navarro-Frenk-White halos, is in good agreement with CDM predictions. Other interesting potentials of unstable CDM are highlighted.

Subject headings: galaxies: halos — galaxies: kinematics and dynamics — galaxies: spiral — dark matter

1. INTRODUCTION

The hierarchical clustering cold dark matter (CDM) scenario for structure formation has been very successful in reproducing most properties of the local and high-redshift Universe. However, some few perceived problems of such models, particularly at small scales, remain open. Numerical studies of collisionless CDM with different initial density fluctuation spectra predict dark halos with steep cusps of the form $\rho(r) \sim \rho_0(r/r_s)^{-\alpha}(1+r/r_s)^{\alpha-3}$, with $\alpha = 1$–1.5, after Navarro, Frenk & White (1996) (NFW profile, hereafter), whereas most of the rotation curves of dwarf galaxies and low surface brightness (LSB) galaxies suggest that their halos have constant density cores (see de Blok, Bosma & McGaugh 2003, and references therein). A possibly related problem is the apparent overprediction of the number of satellites with circular velocities $v_c \sim 10$–30 km s^{-1} within the virialized dark halos of the Milky Way and M31 (Klypin et al. 1999; Moore et al. 1999).

Many astrophysical mechanisms have been suggested to explain these discrepancies. If winds driven by vigorous star formation, bars in forming galaxies and the merger of black holes prove to be inefficient at altering a primordial dark matter cusp (see Gnedin & Zhao 2002, Sellwood 2003 and Read & Gilmore 2003, respectively), one should invoke modifications of the standard CDM scenario. A reduction of the small-scale power by substituting warm dark matter for CDM or by appealing to a model of inflation with broken scale invariance is strongly discouraged by the recent WMAP observations (Bennett et al. 2003; Spergel et al. 2003).

Other alternatives involve altering the fundamental nature of the dark matter itself (Spergel & Steinhardt 2000; Kaplinghat, Knox & Turner 2000; Cen 2001a; Giraud 2001; Arbey et al. 2003, and references therein). All these models have free parameters tunable to fit the observations. For instance, the self-interacting dark matter (SIDM) scenario with a general velocity dependent cross-section per unit mass $\sigma_D/m_D = \sigma_0\,(v_{100}/v)^a$, where $v_{100} = 100$ km s^{-1}, satisfied all the astrophysical requirements for a very narrow range of parameters $\sigma_0 = 0.5$–1 cm^2 g^{-1} and $a = 0.5$–1 (Hennewai & Ostriker 2002; Colín et al. 2002, and references therein). This huge cross-section has made SIDM unappealing to most WIMP and axion theorists.

There are some empirical evidence against SIDM. The central densities of dark matter in mass models with dark matter cores for the elliptical galaxy NGC 4636 are higher than expected in the SIDM scenario with $a \leq 1$ (Loewenstein & Mushotzky 2002). In addition, the halo eccentricity derived for NGC 720 is in good agreement with collisionless dark matter (Buote et al. 2002). Moreover, a significant reduction of the central condensation of satellite subhalos by dark matter collisions, assisted by tidal stripping, would be difficult to reconcile with the large stellar velocity dispersions observed in the dwarf spheroidal galaxies (Stoehr et al. 2002; D'Onghia & Burkert 2003).

The problems discussed above are also faced by other models whose mechanisms to produce a core are based on interactions, either elastic or inelastic, between dark matter particles. The case of annihilating dark matter is an example. This Letter is aimed at investigating a new variant of unstable dark matter, which corresponds to a modification of the proposal by Cen (2001a), to find out whether it is possible to overcome the difficulties associated with previous scenarios but retaining the desired phenomena.

2. UNSTABLE COLD DARK MATTER AND IMPLICATIONS

Although fits to the rotation curves of LSB galaxies favor halos with cores, a fraction of them is acceptably reproduced by divergent NFW profiles as long as they are much less concentrated than typical halos in standard CDM. Cen (2001a) reconsiders the decaying dark matter scenario (e.g., Davis et al. 1981; Turner, Steigman & Krauss 1984; Turner 1985; Dicus & Teplitz 1986; Dekel & Piran 1987; Sciama 1990), and suggests that the concentration of dark matter halos lowers if one-half of the CDM particles decay to relativistic (but nonradiative) particles by redshift $z = 0$. The decaying CDM scenario (DCDM, hereafter) would provide also an explanation for the formation of dwarf spheroidal galaxies with low velocity dispersions (Cen 2001b).

Unfortunately, there are some concerns associated with the DCDM model. First, not all the galaxies can be fitted with a NFW profile varying the concentration parameter, especially dwarf galaxies (e.g., Blais-Ouellette, Amram & Carignan 2001). Secondly, there are empirical evidence to believe that the density of the dark matter halo of the Milky Way at $z = 2$ agrees with the present day halo within 50 kpc (e.g., Zhao 2002, and references therein). The spatial distribution and dynamics of globular clusters give support to this fact. On the other hand, a reduction in the central density of clusters of galaxies and ellipticals by a factor 2^4, as suggested in the DCDM scenario, might be in disagreement with observations (e.g., Arabadjis et al. 2002; Loewenstein & Mushotzky 2002), and could spoil the good agreement found in standard CDM simulations of halo substructure (Stoehr et al. 2002). However, we point out here that the excessive reduction of central dark matter densities in the halos of galaxies and clusters of galaxies can be alleviated if a fraction f of the CDM particles decays to relativistic and nonradiative light particles plus a stable massive particle with a recoiling velocity in the center of mass of the parent particle of, say 30–60 km s^{-1}. If the unstable parent particles are denoted by D and the stable massive particles by D', the decay is given by $D \to D' +$ light particles. We will refer to this model as the unstable CDM model (UCDM). Initially, dark halos are made up of a 'primordial' mixture of particles D and D'. Except for dark matter particles in the outer halo where the escape velocities are smaller than the recoiling velocity, $v_{\rm rc}$, the particles D' created by decay, will remain bound to the galactic halos. Only the light particles that are relativistic will escape from the host halo. Therefore the total mass in a Milky Way halo within 50 kpc would remain approximately constant (see Eq. 2-192 of Binney & Tremaine 1987), consistent with observations. If m_D and $m_{D'}$ are the particle mass of D and D', respectively, a recoiling velocity of a few tens km s^{-1} implies $(m_D - m_{D'})/m_D \lesssim 1$–$2 \times 10^{-4}$. The consequences of UCDM will be discussed in the remainder of this Letter.

2.1. *Formation of Cores in Low-Mass Galaxies*

This section is sought to outline how the profile of dark halos would evolve when CDM particles decay and to elucidate whether a central density core is expected. For simplicity we will assume pure CDM halos and ignore the contribution of the baryonic component. This is a good approximation for galaxies dominated by the dark matter. The implications of the inclusion of the baryonic component is discussed in §2.2. As long as a certain dark halo formed at high redshift and the mean lifetime of dark particles is long enough, the initial halo made up of a mixture of particles D and D', should follow a NFW profile with an inner slope $\alpha \approx 1$–1.5 and a concentration, $c_{\rm i}$, as predicted in cosmological simulations (Navarro et al. 1996, 1997; Power et al. 2003). If f_D is the 'primordial' fraction of particles D and $f_{DD'}(t)$ is the fraction of particles of type D that have decayed at time t, then the fraction of decayed particles is $f(t) = f_D f_{DD'}(t)$. Note that we do not need to employ a fine-tuning between the mean lifetime of D particles, τ, and the present Hubble time $t_{\rm H}$; the case τ much shorter than $t_{\rm H}$ just implies that all particles D have already decayed at the present Hubble time, i.e. $f(t_{\rm H}) = f_{\rm D}$.

The kinetic energy of the daughter particles D' just after decay depends on the angle between the velocity of the parent particle and the recoiling velocity; some of them can even lose energy if, for instance, the recoiling velocity is opposite to the direction of the orbital velocity of the parent particle (except when $v_{\rm rc} \gg \sigma_{\rm t}$, where $\sigma_{\rm t}$ denotes the typical one-dimensional velocity dispersion of halo particles). However, the kinetic energy of the daughter particle, just after decay, averaged over all the possible angles of decay of a parent particle with velocity $\boldsymbol{v}_{\rm D}$ is $\langle E\rangle = m_{\rm D'} v_{\rm D}^2/2 + m_{\rm D'} v_{\rm rc}^2/2$. Therefore, the net increase of energy is $m_{\rm D'} v_{\rm rc}^2/2$, which is independent, on average, of the initial velocity $v_{\rm D}^2$.

The dark matter decay will result in a gradual expansion of the halos. The typical variation in the size of halos comes from virial arguments as follows. A halo with a virial radius $R_{\rm vir,i}$ will expand after the decay of dark particles to a virial radius $R_{\rm vir,f}$ of order of $R_{\rm vir,f}/R_{\rm vir,i} \approx (1 - fx^2/3)^{-1}$, where $x \equiv v_{\rm rc}/\sigma_{\rm t}$. For x small, as occurs in clusters of galaxies (typically $x \approx 0.04$) and giant ellipticals, the effect is negligible, $R_{\rm vir,f} \approx R_{\rm vir,i}[1 + \mathcal{O}(x^2)]$, even if $f = 1$. Accordingly, the profiles of massive galaxies and galaxy clusters should not deviate significantly from the predictions of standard collisionless dark matter in the UCDM hypothesis. However, for objects with low $\sigma_{\rm t}$, as occurs in dwarf galaxies, the inclusion of dark matter decay may have important consequences.

The detailed evolution of the density profile of a certain halo will depend on the distribution function of its particles. The least favorable case for core formation occurs when the halo particles are assumed to be initially on circular orbits because this situation minimizes the kinetic energy in the inner halo. By contrast, in a halo dominated by radial orbits, the circularization of orbits by the decay could partly remove the cusp. In order to quantify the effect of different internal kinematics of the CDM particles, we explore first the case in which most of the dark particles are on nearly circular orbits, followed by an isotropic model.

To illustrate how cores are produced, let us assume that the inner circular velocity $v_{\rm c}(R,t)$ of the initial configuration of the halo, when almost no particle has decayed yet, is given approximately by the power law $v_{\rm c}(R,0) = v_0\,(R/R_0)^{1/2\beta}$, with $\beta > 0$ and v_0 the circular velocity at a galactocentric radius R_0. In the NFW profile, the exponent $1/(2\beta) \approx 1/2$, whereas for a more shallow rotation curve, as the rigid-solid rotation, $1/(2\beta) \approx 1$. Suppose that between $t = 0$ and $t = dt$ a small fraction of particles decays ($f \ll 1$). The daughter particles 'gain' an amount of energy of $\Delta E \approx m_{\rm D'} v_{\rm rc}^2/2$, as said before. If we define the mean radius for non-circular orbits as the radius of a circular orbit with the same energy, it is straightforward to show that the mean radius of the daughter particle, $R_{\rm D'}$, is:

$$R_{\rm D'} = \left(R_{\rm D}^{1/\beta} + \frac{1}{2\beta+1} \left(\frac{v_{\rm rc}}{v_0} \right)^2 R_0^{1/\beta} \right)^{\beta}, \qquad (1)$$

where $R_{\rm D}$ is the mean radius of its parent particle. A *measure* of the expansion of the halo, when the initial orbits are approximately circular, is given by the relative variation in the radius of the particles:

$$\frac{R_{\mathrm{D}'}-R_{\mathrm{D}}}{R_{\mathrm{D}}}=\sum_{i=1}^{\infty}\binom{\beta}{i}\left(\frac{1}{2\beta+1}\right)^{i}\left(\frac{v_{\mathrm{rc}}}{v_0}\right)^{2i}\left(\frac{R_0}{R_{\mathrm{D}}}\right)^{i/\beta}, \tag{2}$$

which goes as $\propto R_{\mathrm{D}}^{-1/\beta}$ in the particular case that $v_{\mathrm{rc}} \lesssim v_0$ and $R_{\mathrm{D}} \gtrsim R_0$, keeping only the leading term. This result suggests that the rotation curve becomes more shallow, once the fraction of mass that decays is not longer negligible at say $t = \delta t$ [1]. Simultaneously, the halo will undergo an adiabatic expansion as a response to the new mass configuration, decreasing the velocity dispersion of the halo particles. This expansion also contributes to the formation of a shallow profile. Inserting the new values of β and v_0 at $t = \delta t$ in Eq. (2), we see that $1/(2\beta)$ always increases in time, concluding that the formation of a core should be a natural consequence in this model. For a more realistic, isotropic orbital distribution, the formation of a core should be even more favorable.

The procedure outlined above allows us to calculate numerically the evolution of a dark halo starting with a NFW profile ($\alpha = 1$) and for different values of v_{rc} (see Fig. 1). For the isotropic models, we have taken advantage of the analytical fit to the distribution function given by Widrow (2002). The initial halo density profile is defined by the concentration parameter c and the rotation velocity V_{200} at radius R_{200}, that is the radius inside which the average overdensity is 200 times the critical density of the Universe. The resulting rotation curves should be regarded as lower limits since the orbits of the daughter particles were assumed to be circular. However, the maximum increment of the rotation velocity by including an isotropic orbital distribution for these particles can be estimated from the differential energy distribution (e.g., Binney & Tremaine 1987) and it is marked with error bars in Fig. 1.

In order to quantify the relevance of the UCDM model, we have recovered the concentration parameter before CDM decay, c_i, for three extreme galaxies: IC 2574, NGC 3274 and Draco. IC 2574 is the prototype dwarf LSB with a conspicuous core (e.g., Blais-Ouellette et al. 2002) whereas NGC 3274 is a LSB galaxy with one of the smallest core-radius, which can be fitted with a NFW profile (de Blok & Bosma 2002). On the other hand, Draco is a dwarf spheroidal (dSph) and presents central dark matter densities (within a radius of ~ 1 kpc) as high as those predicted by standard collisionless CDM models (Kleyna et al. 2001; Lokas 2002; Stoehr et al. 2002; Hayashi et al. 2003). Therefore, these galaxies should bracket the concentration parameters for low-mass galaxies. If the initial concentration parameters for these extreme galaxies, c_i, lie within 1σ scatter of the predictions of standard CDM simulations, it would imply that our model is fully consistent with the observed distribution of concentrations. For the reference values $f = 0.5$ and $v_{\mathrm{rc}} = 40$ km s^{-1}, the observed rotation curves are reproduced well in the isotropic model, starting with a NFW halo with $c_i \approx 13, 26, 20$ and $V_{200,i} \approx 60, 55, 60$ km s^{-1} for IC 2574, NGC 3274 and Draco, respectively. Although cosmological N-body simulations must be done for a fully description of the profile of UCDM halos, these results are promising if one considers that expected values of c_{i} in ΛCDM cosmology have $\sigma(\log c_{\mathrm{i}}) = 0.18$ and a 2σ range from 5 to 25, with an average of around 10 to 15 (NFW; Eke et al. 2001). Since the effects of UCDM are smaller for more massive halos, the present mechanism for core formation is fully consistent with the fact that the best-fitting models of high surface brightness galaxies favor cores with small sizes (Jiménez, Verde & Oh 2003).

2.2. *The Case of Dwarf Spheroidals*

Baryonic effects on the dark matter distribution could affect our estimate of c_i. In particular, if the progenitors of dSph were much more massive than presently, the enhancement of the velocity dispersion of CDM particles by adiabatic compression would have effectively reduced the evacuation of dark matter particles. In the following we will analyze how massive the dSph progenitors should have been to *suppress* significantly the effect of CDM decay. For simplicity, the baryons are assumed to dominate the total mass (dark matter plus baryons) within the optical radius of the galaxy during the phase of CDM decay. The corresponding density profile in this phase is taken to be $\rho \sim R^{-\xi}$, with $\xi \approx 1.8$–2, for consistency with other studies (e.g., Hennawi & Ostriker 2002). The flat circular velocity associated with this configuration will be denoted by v_{c}. Under the approximation that the CDM particles are on circular orbits, a CDM particle initially at a orbital radius R will shrink to a radius $R' = (M/M')R$, where M is the initial mass within R and M' is the final mass within R', during the dissipative collapse of baryons. After decay, the daughter particle will acquire a mean circular radius $R(M/M')\exp\left(v_{\mathrm{rc}}^2/2v_{\mathrm{c}}^2\right)$. Later on, most of the baryonic mass will be ejected probably by supernova explosions or stripped by ram pressure. This mass loss will expand the orbit of the daughter particle to a final radius of $\sim R\exp\left(v_{\mathrm{rc}}^2/2v_{\mathrm{c}}^2\right)$, still assuming that the process evolves adiabatically. The net effect is a self-similar expansion $R \to \lambda R$, with $\lambda \equiv \exp\left(v_{\mathrm{rc}}^2/2v_{\mathrm{c}}^2\right)$ and, therefore, the initial inner velocity profile $v(R) \propto R^{1/2}$, associated with a NFW profile $\rho_{\mathrm{D}} \propto R^{-1}$, is transformed into $v'(R) = \nu v(R)$ with $\nu^2 = 1 - f + f/\lambda^2$. We will employ a variation of $\nu \approx 0.8$, which is compatible with observational uncertainties ($> 25\%$) in the mass determination of dSph (e.g., Kleyna et al. 2001; Lokas 2002), and the fact that, in our model, the mass loss of CDM particles induced by Galactic tidal forces would be significantly suppressed if the progenitor was more compact in the past. For $f = 0.5$, $v_{\mathrm{rc}} = 40$ km s^{-1} and $\nu = 0.8$, a value $v_{\mathrm{c}} \gtrsim 35$ km s^{-1} is required. In terms of the baryonic mass, this implies a mass of gas $\gtrsim 2.5 \times 10^8$ M$_\odot$ for the progenitor of the dSph Fornax with a typical radius of ~ 0.8 kpc, and $\gtrsim 0.4 \times 10^8$ M$_\odot$ for the progenitor of Draco with radial size of ~ 0.15 kpc. As we have assumed circular orbits, these estimates are clearly lower limits. For progenitors with these masses, the decay of UCDM could not have observable effects in their halos. The inferred masses are large, especially for Draco, but are not unreasonable; they are in accordance with different lines of research that point out masses of

---

[1] In the DCDM scenario proposed by Cen (2001a), the byproducts of the decay were assumed to be relativistic. This assumption immediately implies that the adiabatic evolution of the halo is self-similar and hence no core is formed. By contrast, the inclusion of a recoiling velocity breaks down the self-similarity in the evolution.

10^9–10^{10} $M_\odot$ for the progenitors of those dwarf spheroidals with a long period of star formation like Draco and Fornax (Kitayama et al. 2001; Ikuta & Arimoto 2002). In a much more speculative scenario, if the gas mass of dSph is never larger than 10^7 $M_\odot$, dSph could preserve their initial dark halos intact under the assumption that CDM only decays by interactions with baryons/nucleons with a certain cross-section $\sigma_{\rm DN}$.

3. DISCUSSION

The possibility that dark matter decays is being a topic of active work. Theoretical predictions for lifetime of heavy sterile neutrinos and Ultra Heavy particles, as WIMPZILLAs, cover a large range of values, from 10^{-2} to 10^{11} Gyr. We have assumed that UCDM and its byproducts are weakly interacting particles. The possibility that relativistic light particles of the decay are the source of Ultra High Energy Cosmic Rays or responsible for the reionization of the Universe is still open. Even if Ultra Heavy particles were to decay to Standard Model particles, they are not ruled out by present observations (e.g., Ziaeepour 2000).

Besides the formation of density cores, other interesting phenomena associated with UCDM arise. UCDM enhances substructure destruction within galactic halos, and the evacuation of part of the central mass of dark matter in dwarf ellipticals and dwarf spheroidals would lead to an adiabatic expansion of their globular cluster systems, and would partly compensate the drag of dynamical friction, alleviating the problem of the survival of the globular clusters (Lotz et al. 2001, and references therein). Moreover, assuming that UCDM particles are self-interacting with the cross-section found by Hennawi & Ostriker (2002), $\sigma_{DD}/m_D \approx 0.02$ cm^2 g^{-1}, it is feasible to incorporate the growth of supermassive black holes in our model, as suggested by Ostriker (2000). The decay of particles can contribute to feed up the central black hole if they are scattered into the loss-cone, but this contribution is much smaller than the Bondi accretion.

I thank V. Avila-Reese and an anonymous referee for helpful comments.

REFERENCES

Arabadjis, J. S., Bautz, M. W., & Garmire, G. P. 2002, ApJ, 572, 66
Arbey, A., Lesgourgues, J., & Salati, P. 2003, astro-ph/0301533
Bennett, C. L. et al. 2003, astro-ph/0302207
Binney, J., & Tremaine, S. 1987, Galactic Dynamics (Princeton Univ. Press: Princeton)
Blais-Ouellette, S., Amram, P., & Carignan, C. 2001, ApJ, 121, 1952
Buote, D. A., Jeltema, T. E., Canizares, C. R., Garmire, G. P. 2002, ApJ, 577, 183
Cen, R. 2001a, ApJ, 546, L77
Cen, R. 2001b, ApJ, 549, L195
Colín, P., Avila-Reese, V., Valenzuela, O., & Firmani, C. 2002, ApJ, 581, 777
Davis, M., Lecar, M., Pryor, C., & Witten, E. 1981, ApJ, 250, 423
Dekel, A., & Piran, T. 1987, ApJ, 315, L83
de Blok, W. J. G., McGaugh, S. S., & Rubin, V. C. 2001, ApJ, 122, 2396
de Blok, W. J. G., & Bosma, A. 2002, A&A, 385, 816
de Blok, W. J. G., Bosma, A., & McGaugh, S. S. 2003, MNRAS, 340, 657
Dicus, D. A., & Teplitz, V. L. 1986, Phys. Rev. D, 34, 934
D'Onghia, E., & Burkert, A. 2003, ApJ, 586, 12
Eke, V. R., Navarro, J. F., & Steinmetz, M. 2001, ApJ, 554, 114
Giraud, E. 2001, ApJ, 558, L23
Gnedin, O. Y., & Zhao, H. S. 2002, MNRAS, 333, 299
Hayashi, E., Navarro, J. F., Taylor, J. E., Stadel, J., & Quinn, T. 2003, ApJ, 584, 541
Hennawi, J. F., & Ostriker, J. P. 2002, ApJ, 572, 41
Ikuta, C., & Arimoto, N. 2002, A&A, 391, 55
Jiménez, R., Verde, L., & Oh, S. P. 2003, MNRAS, 339, 243
Kaplinghat, M., Knox, L., & Turner, M. S. 2000, Phys. Rev. Lett., 85, 3335
Kitayama, T., Susa, H., Umemura, M., & Ikeuchi, S. 2001, MNRAS, 326, 1353
Kleyna, J. T., Wilkinson, M. I., Evans, N. W., & Gilmore, G. 2001, ApJ, 563, L115
Klypin, A., Kravtsov, A. V., Valenzuela, O., & Prada, F. 1999, ApJ, 522, 82
Loewenstein, M., & Mushotzky, R. 2002, astro-ph/0208090
Lokas, E. L. 2002, MNRAS, 333, 697
Lotz, J. M., Telford, R., Ferguson, H. C., Miller, B. W., Stiavelli, M., & Mack, J. 2001, ApJ, 552, 572
Moore, B., Ghigna, S., Governato, F., Lake, G., Quinn, T., Stadel, J., & Tozzi, P. 1999, ApJ, 524, 19
Navarro, J. F., Frenk, C. S., & White, S. D. M. 1996, ApJ, 462, 563
Navarro, J. F., Frenk, C. S., & White, S. D. M. 1997, ApJ, 490, 493
Ostriker, J. P. 2000, Phys. Rev. Lett., 84, 5258
Power, C., Navarro, J. F., Jenkins, A., Frenk, C. S., White, S. D. M., Springel, V., Stadel, J., & Quinn, T. 2003, MNRAS, 338, 14
Read, J. I., & Gilmore, G. 2003, MNRAS, 339, 949
Sciama, D. W. 1990, ApJ, 364, 549
Sellwood, J. A. 2003, ApJ, 587, 638
Spergel, D. N., & Steinhardt, P. J. 2000, Phys. Rev. Lett., 84, 3760
Spergel, D. N. et al. 2003, astro-ph/0302209
Stoehr, F., White, S. D. M., Tormen, G., Springel, V. 2002, MNRAS, 335, L84
Turner, M. S. 1985, Phys. Rev. D, 31, 1212
Turner, M. S., Steigman, G., Krauss, L. M. 1984, Phys. Rev. Lett., 52, 2090
Widrow, L. M. 2000, ApJS, 131, 39
Zhao, H. S. 2002, MNRAS, 336, 159
Ziaeepour, H. 2000, Astrop. Physics, 16, 101



FIG. 1.— Variations in the shape of the halo contribution to the rotation curve for different values of f and $v_{\rm rc}$, starting with a NFW profile with $c_{\rm i} = 13$ and $V_{200,i} = 60$ km s^{-1} (triple-dotted dashed line) except for the NGC 3274 halo for which $c_{\rm i} = 26$ and $V_{200,i} = 55$ km s^{-1}. Isotropic models are labeled by ISO, while models in which all particles are initially on circular orbits are indicated by CIR. The three solid lines show the rotation curves for $f = 0.5$ and $v_{\rm rc} = 40$ km s^{-1}. The dashed lines correspond to $v_{\rm rc} = 60$ km s^{-1} and $f = 0.5$ (lower dashed curve) and $f = 0.25$ (the two upper dashed curves). The bars account for the maximum effect when including non-circular orbits for the daughter particles for the IC 2574 and NGC 3274 halos.